

Temperature References
Updated :22 April 2009

Climate and Flow

EA

D. Hammond and A.R. Pryce Climate change impacts and water Temperature July 2007 EA Science Report: SC060017/SR ISBN: 978-1-84432-802-4 (#110648)

Fung,C.F, Lopez,A & New,M. Applying probabilistic information to strategic (2009) EA Climate Change Science programme Science report: SC050045 ISBN: 978-1-84911-034-1

Hydrology/Physical

Caissie, D. (2006) The thermal regime of rivers: a review. *Freshwater Biology*, **51, 1389-1406.**

Dymond, J.R.1984: Water temperature change caused by abstraction. Journal of hydraulic engineering 110: 987-991.

Hannah, D.M., Webb, B.W., & Nobilis, F. (2008) River & Stream temperature. Hydrological Processes: 22, 889-901

Langford, T.E. (2001). Thermal discharges and pollution. pp.2933-40 in *Encyclopedia of Ocean Sciences*. Academic Press.

Mackey A.P. & Berrie A.D. (1991) The prediction of water temperatures in chalk streams from air temperatures. *Hydrobiologia*, 210, 183–189.

Ward, J.V (1985) Thermal characteristics of running waters. *Hydrobiologia* 125:31-46

Webb, B.W*. & Langford, T.E.L (2004) Contrasting trends in river water temperatures. Conference on effects of climate change on aquatic organisms. Plymouth.2004

Webb, B.W. & Zhang, Y. (1999) Water temperatures and heat budgets in Dorset chalk water courses. *Hydrological Processes*, **13**, 309-321.

Webb, B.W, Clack,P.D. & Walling, D.E. Water-air temperature relationships in a Devon river system and the role of flow. *Hydrol. Process* 17, 3069-3084 (2003) (#1076025)

Webb, B., & Nobilis, F. (2007) Long-term changes in river temperature and the influence of climatic and hydrological factors. *Hydrological Sciences Journal*, 52, 74-85.

https://connect2.environment-agency.gov.uk/exchange/bilboj.Blandford1.SW/Inbox/River Temperature & Ecology References.EML/1_multipart_xF8FF_2_DM-#1076025-v2-River_Temperature_&_Ecology_References.DOC/C58EA28C-18C0-4a97-9AF2-036E93DDAFB3/DanaInfo=webmail.ea.gov,SSL+DM-#1076025-v2-River_Temperature_&_Ecology_References.DOC?attach=1

Webb B.W, Hannah, D.M. Moore R D, Brown L.E & Noblis, F (2008) Recent advances in stream and river Temperature research. *Hydrol. Process.*(2008) (#1100795)

Ecology- invertebrates

Beche, L.A., McElravy, E.P. & Resh, V.H. (2006) Long-term seasonal variation in the biological traits of benthic-macroinvertebrates in two Mediterranean-climate streams in California, USA. *Freshwater Biology*, **51**, 56-75.

Beche, L.A. & Resh, V.H. (2007) Short-term climatic trends affect the temporal variability of macroinvertebrates in California 'Mediterranean' streams. *Freshwater Biology*, **52**, 2317-2339.

Bonada, N., Doledec, S. & Statzner, B. (2007) Taxonomic and biological trait differences of stream macroinvertebrate communities between Mediterranean and temperate regions: implications for future climatic scenarios. *Global Change Biology*, **13**, 1658-1671.

Bonada, N., Rieradevall, M. & Prat, N. (2007) Macroinvertebrate community structure and biological traits related to flow permanence in a Mediterranean river network. *Hydrobiologia*, **589**, 91-106.

Boulton, A.J. (2003) Parallels and contrasts in the effects of drought on stream macroinvertebrate assemblages. *Freshwater Biology*, **48**, 1173-1185.

Bradley, D.C. & Ormerod, S.J. (2002a) Evaluating the precision of kicksampling in upland streams for assessments of long-term change: the effects of sampling effort, habitat and rarity. *Archiv Für Hydrobiologie*, **155**, 199–221.

Briers, R.A., Gee, J.H.R. & Geoghegan, R. (2004) Effects of the North Atlantic Oscillation on growth and phenology of stream insects. *Ecography*, **27**, 811-817.

Clark, E., Webb, B.W. & Ladle, M. (1999) Microthermal gradients and ecological implications in Dorset rivers. *Hydrological Processes*, **13, 423-438.**

Daufresne, M., Bady, P. & Frugé, J. (2007) Impacts of global changes and extreme hydroclimatic events on macroinvertebrate community structures in the French Rhone River. *Oecologia*, **151**.

Daufresne, M., Roger, M.C., Capra, H. & Lamouroux, N. (2004) Long-term changes within the invertebrate and fish communities of the Upper Rhone River: effects of climatic factors. *Global Change Biology*, **10**, 124-140.

Daufresne, M., Bady, P. & Frugé, J. (2007) Impacts of global changes and extreme hydroclimatic events on macroinvertebrate community structures in the French Rhone River. *Oecologia*, **151.**

https://connect2.environment-agency.gov.uk/exchange/bilboj.Blandford1.SW/Inbox/River Temperature & Ecology References.EML/1_multipart_xF8FF_2_DM-#1076025-v2-River_Temperature_&_Ecology_References.DOC/C58EA28C-18C0-4a97-9AF2-036E93DDAFB3/DanaInfo=webmail.ea.gov,SSL+DM-#1076025-v2-River_Temperature_&_Ecology_References.DOC?attach=1

Dewson, Z.S., James, A.B.W. & Death, R.G. (2007) A review of the consequences of decreased flow for instream habitat and macroinvertebrates. *Journal of the North American Benthological Society*, **26**, 401-415.

Durance I & Ormerod, S.J. Evaluating climatic effects on aquatic invertebrates in southern English rivers. February 2008. EA Science report SR. (#935409)

Hall, C.A.S., Stanford, J.A. & Hauer, F.R. (1992). The distribution and abundance of organisms as a consequence of energy balances along multiple environmental gradients. *Oikos* 65: 377-390.

Harper, M.P., & Peckarsky, B.L. (2006). Emergence cues of a mayfly in a high-altitude stream ecosystem: potential response to climate change. *Ecological Applications*, 16, 612-621.

Hauer, F.R., Stanford, J.A., Giersch, J.J., and Lowe, W.H (2000) Distribution and abundance patterns of macroinvertebrates in a mountain stream: an analysis along multiple environmental gradients Verh. Internat. Verein. Limnol. 27: 1485-1488.

Hawkins C.P., Hogue J.N., Decker L.M. & Feminella J.W. (1997) Channel morphology, water temperature, and assemblage structure of stream insects. *Journal of the North American Benthological Society*, 16, 728–749.

Hogg, I.D, D.D. Williams, J.M. Eadie, and S.A. Butt (1995). The consequences of global warming for stream invertebrates: a field simulation. *Journal of Thermal Biology* 20: 199-206.

Hogg, I.D. & Williams, D.D. (1996) Response of stream invertebrates to a global-warming thermal regime: An ecosystem-level manipulation. *Ecology*, **77**, 395-407.

Hynes, H.B.N (1970). *The ecology of running waters*. Liverpool University Press. Liverpool.

Lake, P.S. (2000) Disturbance, patchiness, and diversity in streams. *Journal of the North American Benthological Society*, **19**, 573-592.

Lytle, D.A. & Poff, N.L. (2004) Adaptation to natural flow regimes. *Trends in Ecology & Evolution*, **19**, 94-100.

Langford, T.E. (1970). The temperature of a British River upstream and downstream of the heated discharge from a power station. *Hydrobiologia* **35** (3-4), 353-375.

Langford, T.E. (1972). A comparative assessment of thermal effects in some British and North American rivers. In: *River Ecology and Man*. R.T. Oglesby, C.A., Carlson & J.A. McCann (Eds.) pp. 315-351. Academic Press. New York. London.

https://connect2.environment-agency.gov.uk/exchange/bilboj.Blandford1.SW/Inbox/River Temperature & Ecology References.EML/1_multipart_xF8FF_2_DM-#1076025-v2-River_Temperature_&_Ecology_References.DOC/C58EA28C-18C0-4a97-9AF2-036E93DDAFB3/DanaInfo=webmail.ea.gov,SSL+DM-#1076025-v2-River_Temperature_&_Ecology_References.DOC?attach=1

Langford, T.E. (1974). Ecology and cooling water from power stations – A review of recent biological research in Britain. In: *The Proceedings of the World Energy Conference, Detroit. Paper 2.2.4.* WEC. London.

Langford, T.E. (1975). The distribution, abundance and life-histories of stoneflies (Plecoptera) And mayflies (Ephemeroptera) in a British river warmed by the cooling water from a power station. *Hydrobiologia* **38** (2), 339-377.

Langford, T.E. (1975). The emergence of insects from a British river warmed by power station cooling water. Part 11. The emergence patterns of some species of Ephemeroptera, Trichoptera and Megaloptera in relation into water temperature and river flow upstream and downstream of the cooling water outfalls. *Hydrobiologia* **47** (1), 91-133.

Langford, T.E. & Aston, R.J. (1972). The ecology of some British rivers in relation to warm water discharges from power stations. *Proceedings of the Royal Society, London* **181**, 45-47.

Langford, T.E. & Bray, E.S. (1969). The distribution of Plecoptera and Ephemeroptera in a Lowland Region of Britain (Lincolnshire). *Hydrobiologia* **34** (2), 243-271.

Langford, T.E. & Daffern, J.R. (1975). The emergence of insects from a British river warmed by power station cooling water. Part 1. The use and performance of insect emergence traps in a large spate river and the effects of various factors on total catches upstream and downstream of the cooling water outfalls. *Hydrobiologia* **46** (1), 71-114.

Langford, T.E., James, N.P. & Fleming, J.M. (1978). Power station studies. UK. *Underwater Telemetry Newsletter* **8** (1). July. Environmental Sciences Division. Oak Ridge Laboratory.

Langford, T.E.L. and Langford, J. 2002. *Effects of water use at Power Stations on aquatic ecosystems and methods of assessment.* A generic review of current information and methodologies in relation to future regulations, 2002. Pisces Conservation Ltd. (Client: TXU – Europe Plc).

Mouthon, J. & Daufresen, M. (2006). Effects of the 2003 heatwave and climatic warming on mollusc communities of the Saone: a large lowland river and of its two main tributaries (France). *Global Change Biology*, 12, 441-449.

https://connect2.environment-agency.gov.uk/exchange/bilboj.Blandford1.SW/Inbox/River Temperature & Ecology References.EML/1_multipart_xF8FF_2_DM-#1076025-v2-River_Temperature_&_Ecology_References.DOC/C58EA28C-18C0-4a97-9AF2-036E93DDAFB3/DanaInfo=webmail.ea.gov,SSL+DM-#1076025-v2-River_Temperature_&_Ecology_References.DOC?attach=1

Pritchard, G., Harder, L.D. & Mutch, R.A. (1996) Development of aquatic insect eggs in relation to temperature and strategies for dealing with different thermal environments. *Biological Journal of the Linnean Society*, **58**, 221-244.

Quinn, J.M., Steele, G.L., Hickey, C.W., & Vickers, M.L. (1994). Upper tolerances of twelve New Zealand stream invertebrate species. *New Zealand Journal of Marine and Freshwater Research*. Vol. 28: 391-397

Rostgaard, S. & Jacobsen, D. (2005) Respiration rate of stream insects measured in situ along a large altitude range. *Hydrobiologia*, **549**, 79-98.

Townsend CR, Hildrew AG, Francs JE (1983): Community structure in some southern English streams: the influence of physiochemical factors. *Freshwater Biology* 13: 521-544.

Townsend CR, Scarbrook, M.R., Doledec, S. (1997) The intermediate disturbance hypothesis, refugia, and biodiversity in streams. *Limnology and Oceanography*, 42, 938-949.

Vannote, R.L., Sweeney, B.W (1980). Geographic analysis of thermal equilibria – a conceptual-model for evaluating the effect of natural and modified thermal regimes on aquatic insect communities. *American Naturalist*, 115, 667-695.

Vannote, R.L., Minshall, GW., Cummins, K.W., Sedell, J.R., Cushing, C.E. (1980) The river continuum concept. *Canadian Journal of Fisheries and Aquatic Sciences*, 37, 130-137.

Williams, D.D., and N.E. Williams. 1993. The upstream/downstream movement paradox of lotic invertebrates: quantitative evidence from a Welsh mountain stream. *Freshwater Biology* 30: 199-218.

Wood, P.J., Hannah, D.M., Agnew, M.D. & Petts, G.E. (2001) Scales of hydroecological variability within a groundwater-dominated stream. *Regulated Rivers-Research & Management*, **17**, 347-367.

Woodward, G., Jones, J.I. & Hildrew, A.G. (2002) Community persistence in Broadstone Stream (UK) over three decades. *Freshwater Biology*, 47, 1419–1435.

Wright, J.F., Clarke, R.T., Gunn, R.J.M., Kneebone, N.T. & Davy-Bowker, J. (2004) Impact of major changes in flow regime on the macroinvertebrate assemblages of four chalk stream sites, 1997-2001. *River Research and Applications*, **20**, 775-794.

Fish

Langford, T.E. (2006).** Temperature and Fish Conservation. Institute of Fisheries Management Conference. Minehead.

https://connect2.environment-agency.gov.uk/exchange/bilboj.Blandford1.SW/Inbox/River Temperature & Ecology References.EML/1_multipart_xF8FF_2_DM-#1076025-v2-River_Temperature_&_Ecology_References.DOC/C58EA28C-18C0-4a97-9AF2-036E93DDAFB3/DanaInfo=webmail.ea.gov,SSL+DM-#1076025-v2-River_Temperature_&_Ecology_References.DOC?attach=1

Solomon D.J & Lightfoot G, Thermal Biology of Brown trout and atlantic salmon.
August 2008EA Science report SR Published by: Environment Agency
ISBN: 978-1-84432-932-8 (#1030148)

Webb, B.W & Walsh, A.J (2004) Changing UK river temperatures and their impact on fish populations. Hydrology; Science and practice for the 21st Century Volume II (#1100777)

Turnpenny, A.W.H., Langford, T.E. & Aston, R.J. (1985). Power stations and fish. *CEGB Research.* 17 (April) 27-39.

If highlighted in red or # number– Wessex water already hold.

https://connect2.environment-agency.gov.uk/exchange/bilboj.Blandford1.SW/Inbox/River Temperature & Ecology References.EML/1_multipart_xF8FF_2_DM-#1076025-v2-River_Temperature_&_Ecology_References.DOC/C58EA28C-18C0-4a97-9AF2-036E93DDAFB3/DanaInfo=webmail.ea.gov,SSL+DM-#1076025-v2-River_Temperature_&_Ecology_References.DOC?attach=1